Atmospheric wave observations at Arecibo

C. J. ZAMLUTTI

Instituto de Pesquisas Espaciais (INPE), São José dos Campos, São Paulo, Brazil

(Received 5 August 1974)

Abstract—Multiple pulse incoherent scatter experiments, observing *E*-region altitudes with height resolution better than 2.4 km, have been carried out at Arecibo from August 1972 to January 1973. Neutral temperature, ion–neutral collision frequency and drift velocities were obtained. The results showed an oscillatory behavior of the neutral atmosphere. Evidences of non-linearities of the atmospheric oscillations are explored. They help to determine the period of the tidal mode consistent with the semi-diurnal tide predicted by the theory.

1. INTRODUCTION

INCOHERENT scatter measurements carried out at *E*-layer altitudes (90–140 km) are of particular interest in the study of the neutral atmosphere. This is due to the fact that, at these altitudes, neutral and ionized particles are under thermal equilibrium, the collision frequency between ions and neutrals may be determined from the measured autocorrelation functions (or its Fourier transform, the power spectrum), and that winds constitute the main component of the drift velocities. As a result, neutral temperature, neutral particles concentration and wind velocities can be determined from *E*-layer incoherent scatter measurements.

E-layer drift velocity measurements, recently improved at Arecibo (ZAMLUTTI, 1973; ZAMLUTTI and FARLEY, 1974) make available now enough information for more detailed conclusions to be drawn about the dynamics of the neutral atmosphere.

Experiments using the technique of multiple pulses (ZAMLUTTI and FARLEY, 1974) were carried out from September 1972, to January 1973. Two pulse schemes were used: one with seven pulses, and the other with five. The seven-pulse experiments have a height resolution of better than 1.35 km, and yield data on electron density, ion temperature, and ion–neutral collision frequency. The five pulse experiments have a height resolution of better than 2.4 km. These latter experiments sacrifice the height resolution in order to obtain drift velocities as well as the other parameters.

The results of our observations permitted the determination of the dominant periods of oscillation in ion temperature, ion–neutral collision frequency and wind velocities. These periods showed the predominance of one tidal model and two gravity wave modes. The energy of the tidal mode was observed to be much larger than that of the gravity wave modes.

The observed oscillations cannot be explained in terms of the linear theories of gravity and tidal waves (HINES, 1960; LINDZEN, 1970; LINDZEN and BLAKE, 1971; LINDZEN, 1971). The evidences of non-linearities are explored here. It appears more realistic to interpret the observed oscillations as being due to non-linear waves (CHO, 1972).

2. DATA SOURCE

Incoherent scatter multiple pulse experiments, were carried out, at Arecibo, from September 1972 to January 1973. Two pulse schemes were used: one with seven pulses and altitude resolution of better than 1.35 km, the other with five pulses and altitude resolution of better than 2.4 km. Details about these experiments are given by ZAMLUTTI and FARLEY (1974). The experiments yielded data on electron density, ion temperature, ion-neutral collision frequency and drift velocity.

The observations were required to last more than 12 hr to resolve the tidal modes of oscillation. Similarly an integration time of better than 30 minutes has been imposed to resolve the gravity wave modes.

Neutral temperature and ion neutral collision frequency were obtained using the analysis method described by ZAMLUTTI (1973). It was assumed that the ion composition was 50 % NO^+ and 50 % O_2^+ which yields a mean ion mass of 31 a.m.u. Thermal equilibrium was also admitted to hold at E -layer heights.

Drift velocity data were obtained with the analysis method described by ZAMLUTTI (1973). Wind velocity, as shown by BERNARD and SPIZZICHINO (1971), can be assumed to be equal to the drift velocity at E -layer heights. To determine the vertical and northward horizontal component of winds, the antenna beam was tilted with respect to the zenith and pointed alternatively to the north and to the south. For our purposes here, however, only line-of-sight velocities really matter.

3. ANALYSIS OF DATA

The constituent periodicities of the diurnal oscillations observed in the ion temperature and in the ion neutral collision frequency were determined using a variant of Prony's method (HILDEBRAND, 1956). Basically the technique is very simple: it consists of finding the angular frequencies, ω_i ($i = 1, 2, 3, \dots, N$), that best describe the oscillatory behavior of a function, $f(x)$, when it is developed as:

$$f(x) = A_0 + \sum_{i=1}^N A_i \cos \omega_i x + B_i \sin \omega_i x$$

where N is the number of modes that we want to consider. A least squares fitting procedure yields the amplitudes A_i , B_i , once the angular frequencies ω_i , are known.

The drift velocity data were analyzed using the correlation technique (BERNARD and SPIZZICHINO, 1971). To observe the effect of the magnetic field a separate analysis has been performed on the data obtained with the antenna beam directed to the north (angle between the direction of propagation and the magnetic field $\alpha = 52^\circ$) and the data with the antenna beam directed to the south ($\alpha = 28^\circ$).

4. RESULTS

The distribution of dominant periods observed in ion-neutral collision frequency and ion temperature data are shown in the histograms of Figs. 1 and 2. A good agreement of Fig. 1 with Fig. 2 can be observed for the distribution of short periods (around 2.4 hr and 4.0 hr). For the long periods with mean value around 13 hr the results obtained with ion-neutral collision frequency data differ from the

Fig. 1. Histogram showing the number of occurrences of the periods of oscillations determined from the ion-neutral collision frequency data.

Fig. 2. Histogram showing the number of occurrences of the periods of oscillations determined from the ion temperature data.

ones obtained with ion temperature data. It can also be observed that the long periods are widely distributed around their mean value.

The apparent disagreement between Figs. 1 and 2 as far as the long periods of oscillation are concerned, can be easily explained by the presence of sporadic *E*-layers (often observed at Arecibo during our experiments). As mentioned before, our data has been analyzed with the assumption of an ion mass of 31 a.m.u. This assumption is certainly always valid when poor altitude resolution (more than 10 km) is used. In this case the effect of any mild sporadic *E*-layer is average out. When good altitude resolution is used (better than 3 km), care should be taken with this assumption. In this case of a highly dense sporadic *E*-layer reasonable results for ion temperature and ion-neutral collision frequency can only be achieved if the actual ion composition is taken into account (ZAMLUTTI, 1973; ZAMLUTTI and FARLEY, 1974).

It has been verified that for highly dense sporadic *E*-layers if we neglect the actual composition, the data analysis program yields poor results for the ion-neutral collision frequency, while still providing tolerable results for the ion temperature (no more than 20 per cent off). This means that for mild sporadic

E-layers (like the ones present during our experiments) we may expect the ion-neutral collision frequency data to be less reliable than the ion temperature data. For this reason we better trust our results of ion temperature and therefore further discussions will mostly be based on them.

A quantitative appreciation for the distribution of periods shown in Fig. 2 is presented in Table 1. The ratio between the standard deviation, σ , and the mean value, μ , was used to give an idea of how the periods are distributed around their mean value.

Table 1.

$\mu(\text{hr})$	$ \sigma/\mu $
2.4	0.1
3.9	0.1
13.1	0.2

WAND (1969) and WAND and PERKINS (1970) analyzed ion temperature data from Arecibo and found, as we did, a widely spread distribution of the dominant tidal mode. The dominant periods in their case ranged from 7–13 hr.

BERNARD and SPIZZICHINO (1971) also studied the tidal modes of oscillation of the neutral atmosphere. Using a correlation analysis they averaged 36 days of data. As a result they found, from ion temperature data, an average period of 10.5 hr with 2 hr standard deviation, and from wind velocity data an average period of 12 hr with 1 hr standard deviation.

Table 1 shows that the gravity wave modes are less spread around their mean value than the tidal mode is. This is simple to understand. In the 12-hr data, several periods of gravity wave oscillations are observable. For the same experiment duration only one complete period of the tidal oscillation can be observed. This latter is, thus, not as well determined in the analysis to find the constituent periodicities.

The profile for the amplitude of the oscillations, observed in the ion temperature data on December 1972 is shown in Fig. 3. One can note that most of the energy of the oscillations is concentrated in the tidal mode. The amplitude of the oscillation in ion temperature has been observed not to exceed the value of 25 per cent of its diurnal average.

The linear theory of neutral atmosphere oscillations predicts amplitude profiles which grow monotonically with height in the range 90–130 km. This happens even for relatively large amplitudes (LINDZEN, 1971).

The amplitude profile in Fig. 3 exhibits a wavelike structure with some phase coherence between modes. This strongly suggests the presence of interactions predicated by non-linear theories (CHO, 1972).

Our wind velocity data yield more reliable results since it is independent of the particular ion composition assumed to determine the ion temperature and ion-neutral collision frequency. However, since we have much less data on wind velocity it will be used only to complement our discussion.

Fig. 3. Amplitude of the oscillatory components present in the ion temperature data obtained on 4 December 1972.

Fig. 4. Normalized autocorrelation functions, $\langle v(t, h)v(t + \tau, h) \rangle$, computed with line-of-sight drift velocities, $v(t, h)$, observed on 4 December 1972. The left hand side corresponds to the data obtained with the antenna pointing northward. The right side corresponds to the data with the antenna pointing south.

The cross correlation technique (BERNARD and SPIZZICHINO, 1971) applied to the wind velocity data obtained on 4 December 1972 gave the results shown in Figs. 4 and 5. As mentioned before, we used line-of-sight velocity and kept separate the data obtained with the antenna pointing to the north ($\alpha = 52^\circ$) from that obtained with the antenna pointing to the south ($\alpha = 28^\circ$). Up to 110 km the autocorrelation functions for the data with the beam pointing north do not differ

Fig. 5. Normalized cross-correlation functions, $\langle v(t, h)v(t + \tau, h + \Delta h) \rangle$, computed with line-of-sight drift velocities, $v(t, h)$, observed on 4 December 1972. The left hand side corresponds to the data obtained with the antenna pointing northward. The right hand side corresponds to the data with the antenna pointing south.

significantly from those with the beam pointing to the south. Above 110 km differences can be observed. The drift velocity for the antenna pointing to the north may be affected by an $E \times B$ drift if large electric fields are present (EVANS, 1972) while the neutral wind will continue to be the dominant component in the velocity for the beam pointing south. The predominance of a tidal mode can be noted in Figs. 4 and 5. The distribution of periods yielded a mean value between 12–13 hr in agreement with the results from ion temperature data.

Computing the vertical and northward horizontal component of the velocity we found that the horizontal velocity may be as large as 100 m/sec. This means that the ratio, ε , given by:

$$\varepsilon = (\text{wind velocity}) \times (\text{sound velocity})^{-1}$$

does not satisfy the condition $\varepsilon \ll 1$. Under these circumstances non-linear terms become important in gravity wave theory (CHO, 1972).

In order to observe the effect of non-linearities we considered the distribution of periods observed in ion temperature data with the amplitude of oscillations that satisfy the relation: $\Delta T_n/T_n \leq 0.13$. These oscillations can be assumed to be linear. Figure 6 shows the distribution of such periods. It may be noted that the distribution for the gravity wave modes was not drastically affected. This excludes

Fig. 6. Histogram showing the periods observed in the ion temperature oscillations for a relative amplitude of $\Delta T_n/T_n < 0.13$.

the possibility of these observed periods being the result of the non-linearities of the tidal oscillation. The distribution of the tidal mode was more affected, yielding a new mean value of 12 hr (in agreement with the semi-diurnal period predicted by the linear theory). The distribution around the mean value was only slightly affected and the ratio $|\sigma/\mu|$ became 0.15.

This last result makes it easy to understand our preceding ones. Under conditions of large amplitude oscillations non-linearities become important. Assuming these oscillations are periodic a pseudo-period will be determined from them. This explains why the values found for the semi-diurnal tide were not equal 12 hr.

Other evidence in favor of non-linear effects is seen in Fig. 7, where the diurnal average profile for 7 November 1972 and 18 January 1973 are shown. On 7 November the relative amplitude of the oscillations observed for the neutral temperature was smaller than 10 per cent for all observed heights. On 18 January, the relative amplitude of the oscillations was larger than 20 per cent for all observed heights. The diurnal average neutral temperature profile for this last day

Fig. 7. Diurnal average neutral temperature profiles obtained on 7 November 1972 and 18 January 1973. About 12 hr of data were analyzed.

drastically differs from the monotonic profiles predicted by the hydrostatic equilibrium assumption (which is a reasonable one for small amplitude oscillations) used to predict temperature profiles (JACCHIA, 1970).

5. CONCLUSIONS

Three atmospheric parameters were determined using the incoherent scatter technique. These parameters presented an oscillatory behavior. The methods to determine the periodicity of the data revealed the presence of three dominant modes: 2.4 hr, 3.9 hr and 13.1 hr. Internal consistency of the periods determined from the different parameters proved to be satisfactory. The tidal oscillation prevailed over the others, with most of the energy of the oscillation being concentrated on it.

From our observations we concluded that wind velocity data is more appropriate to determine atmospheric wave periods than ion temperature and ion-neutral collision frequency data. This follows from the fact that for monostatic measurements drift velocity derivation can be made practically insensitive to the assumption of composition necessary to determine the other parameters (ZAMLUTTI, 1973).

The amplitude profile of the temperature oscillations for the range 100–120 km revealed itself as an oscillatory behavior. This suggests that the oscillations are not linear. In fact we observed that the condition for linearity was not satisfied. Restricting our analysis to the neutral temperature oscillations with relative amplitude smaller than 13 per cent, we observed some changes in the distribution of the tidal oscillation periods. The most important was that the mean value period turned out to be 12 hr (instead of the 13.1 hr previously determined) which agrees with that predicted by the linear theories.

The wide distribution around the mean value observed for the tidal oscillations when compared to the gravity wave distribution is easy to explain. In 12-hr experiments only one entire period of the tidal mode can be observed against at least three of the gravity wave modes. Thus the uncertainty in the tidal mode mean value should be roughly twice as much as that of the gravity wave modes. This agrees with what we observed.

Acknowledgements—The cooperation of Prof. D. T. FARLEY and Dr. H. C. CARLSON is gratefully acknowledged. It should be emphasized helpful discussion with Prof. T. HAGFORS and the help of the staff of the National Astronomy and Ionosphere Center at Arecibo. Special mention is due to Mr. P. WATKINS and Dr. G. GIERALTOWSKI for their help. Support for this work came from Instituto de Pesquisas Espaciais (INPE—Brazil) and the National Aeronautics and Space Administration (NASA) under a joint international fellowship program.

REFERENCES

BERNARD R. A. and SPIZZICHINO A.	1971	<i>J. atmos. terr. Phys.</i> 83 , 1345.
EVANS J. V.	1972	<i>J. geophys. Res.</i> 77 , 2341.
HILDEBRAND F. B.	1956	<i>Introduction to Numerical Analysis</i> . McGraw-Hill, New York.
HINES C. O.	1960	<i>Can. J. Phys.</i> 38 , 1441.
LINDZEN R. S.	1970	<i>Geophys. Fluid Dyn.</i> 1 , 303.
LINDZEN R. S. and D. BLAKE	1971	<i>Geophys. Fluid Dyn.</i> 2 , 31.

LINDZEN R. S.	1971	<i>Geophys. Fluid Dyn.</i> 2 , 89.
WAND R. H.	1969	<i>J. geophys. Res.</i> 74 , 5688.
WAND R. H. and PERKINS F. W.	1970	<i>J. atmos. terr. Phys.</i> 32 , 1921.

Reference is also made to the following unpublished material:

CHO H. R.	1972	Tech. Rept. 48 , Univ. of Illinois.
JACCHIA L. G.	1970	Special Rept. 313, Smithson. Astrophys. Obs.
ZAMLUTTI C. J. and FARLEY D. T.	1974	Incoherent Scatter Multiple Pulse Measurements (to be published).
ZAMLUTTI C. J.	1973	Incoherent Scatter Studies of the Lower Ionosphere at Arecibo using Multiple-pulse Techniques, Ph.D. Thesis, School of Electrical Engineering, Cornell Univ.