



XIX CBMET

CONGRESSO BRASILEIRO DE METEOROLOGIA

JOÃO PESSOA PB | 07 A 11 DE NOVEMBRO DE 2016

METEOROLOGIA: TEMPO, ÁGUA E ENERGIA



THE STORM TRACK OVER SOUTH AMERICA: A COMPARISON BETWEEN THE CMIP5 HADGEM2-ES HISTORICAL SIMULATION AND THE ERA-INTERIM REANALYSIS FOR THE RECENT PAST CLIMATE

P. E. Dias da Silva¹, M. M. Coutinho² and K. I. Hodges³

¹ Center for Earth System Science (CCST), National Institute for Space Research (INPE), São José dos Campos, Brazil, philipp.edson@inpe.br,

² Department of Aerospace Science and Technology (DCTA), São José dos Campos, Brazil, marianemmc@iae.cta.br

³ Department of Meteorology, University of Reading, Berkshire, England, United Kingdom, k.i.hodges@reading.ac.uk

1. INTRODUCTION

Cyclones play an important role in the energy transfer between the equator and polar regions. Preferred regions of cyclones propagation are called Storm Tracks (ST). The ST are important in modulating weather and climate in South America (SA), some recent studies have shown that they may be being affected by climate change [1]. This study aims to evaluate the ability of the Hadley Centre Global Environment Model version 2, with its Earth system configuration (HadGEM2-ES) [2], in simulating the ST in the recent past climate. HadGEM2-ES was one of the models used to produce climate simulations for the latest report of the Intergovernmental Panel on Climate Change (IPCC-AR5) [3].

2. METODOLOGY

Storm Tracks are studied for the recent past climate, using an objective feature tracking algorithm (Track) [4] applied to extratropical cyclones. The aim is to provide an indication of the reliability of using HadGEM2-ES over South America to investigate future changes in ST. Track was applied to 27 years (1979-2005) of relative vorticity data at 850hPa from the HadGEM2-ES historical experiment and the ERA-Interim reanalysis [5]. Statistical analysis of the track, genesis and lysis densities were explored to identify key patterns and possible trends in the simulations.

3. RESULTS

The track density statistic for the reanalysis data (Fig.1.a, b) shows the main patterns seen in the literature, such as the ST being closer (more distant) to the pole in austral summer (winter). Regions of genesis are found at the southeast of SA (Fig.1.e, f) and regions of lysis to the west of the Andes (not shown). The track (Fig.1.c, d) and genesis (Fig.1.g, h) densities for HadGEM2-ES shows similar patterns to those of the reanalysis, only slightly displaced.




Figure 1. Lower-tropospheric seasonal cycle of the SH storm track based on ξ_{850} track density for ERA-Interim in a) DJF and b) JJA and for HadGEM2-ES in c) DJF and d) JJA. For e), f), g) and h), same as in the track density, but for genesis density.

The difference between the track densities provided by HadGEM2-ES and ERA-Interim reanalysis (Fig.2.a, b) indicates a positive bias in an area extending from Paraguay and its neighborhood to the southwest of the state of Minas Gerais and another one in an area to the east of the province of Santa Cruz in Argentina. On the other hand, negative biases are observed in the vicinity of Buenos Aires and also at the southern end of the SA. Hence, the analysis shows a zonal bias, positive/negative in similar positions for all seasons, extending from the southeastern Brazil to the south of SA (~50°S).

In the Antarctic Peninsula region the bias is negative in the austral summer/winter and positive in the austral fall/spring (not shown), which shows that the model has difficulties in representing the ST here.




Figure 2 – Lower-tropospheric biases (HadGEM2-ES vs. ERA-Interim) of the SH storm track based on ξ_{850} for track density in a) DJF and b) JJA, genesis density in c) DJF and d) JJA and lysis density in e) DJF and f) JJA.

The genesis density (Fig.2.c, d) is well simulated by the model when compared with the reanalysis, especially for the area of cyclogenesis to the east of southeastern Brazil during the summer. However, the model tends to displace the genesis slightly to the north, which explains the zonal bias observed in all seasons. Also, the model underestimates cyclogenesis in the region to the east of northern Argentina, especially in DJF. Finally, the model represents the lysis density (Fig.2.e, f) similarly to the reanalysis, especially in the most important region to the west of the Andes. However, the model tends to overestimate the values and displaces the lysis areas slightly to the north, except in the austral winter.

4. CONCLUSION

Overall, results show SA cyclones correspond well between the HadGEM2-ES and the ERA-Interim reanalysis, with some regions slightly displaced in the variables analyzed. However, a zonal bias is observed in the track density around the pole associated with an underestimation of the cyclones by HadGEM2-ES.

5. REFERENCES

- [1] IPCC. *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel of Climate Change*. Cambridge University Press, v. 27, 2014. p. 1499-1566.
- [2] JONES, C. D.; HUGHES, J. K.; BELLOUIN, N. et al. *The HadGEM2-ES implementation of CMIP5 centennial simulations*. *Geosci. Model Dev.* 2011. p. 689–763.
- [3] IPCC. Working Group I Contribution to the IPCC Fifth Assessment Report - Summary for Policymakers. In: *Climate Change 2013: The Physical Science Basis*. [s.n.], 2013. p. 1–36.
- [4] HODGES, K. I. *A General Method for Tracking Analysis and Its Application to Meteorological Data*. *Monthly Weather Review*, v. 122, n. 11, p. 2573–2586, 1994. ISSN 0027-0644.
- [5] DEE, D. P.; UPPALA, S. M.; SIMMONS et al. *The ERA-Interim reanalysis: Configuration and performance of the data assimilation system*. *Quarterly Journal of the Royal Meteorological Society*, v. 137, p. 553–597, 2011. ISSN 00359009.

ACKNOWLEDGMENTS

The authors would like to acknowledge the Coordination of Higher Education Personnel (CAPES) - Postgraduate Support Program (PROAP) - and the Postgraduate in Earth System Science (PGCST) for financial support. The ERA-Interim data has been produced by ECMWF and the HadGEM2-ES has been provided by MetOffice as part of a special project with INPE-Brazil.